Version of Specification with Markings to Show Changes

Page 1, paragraphs 1-2

continuation application is οf PCT a International Application No. PCT/US99/22402, filed September 24, 1999, designating the United States of America, which [is claiming] claims the priority of U.S. Provisional Application No. 60/101,693, filed September 25, 1998, the contents of which are hereby incorporated by reference into the present application. []

[

The present application <u>also</u> claims the benefit of U.S. Provisional Application No. 60/101,825, filed September 25, 1998, [and U.S. Provisional Application No. 60/101,693, filed September 25, 1998,] which [are] <u>is</u> incorporated by reference herein.

Page 7, paragraph 5, lines 24-27

Figures 1a-1, 1a-2 and 1a-3 provide[s] the distribution of alanine in the molecular markers (TV-markers) described in Table 1. The amino acid position is defined by the X-axis. The presence of an alanine is indicated by a vertical bar at the indicated amino acid position.

Page 7, paragraph 5, lines 24-27

Figures 1b-1, 1b-2 and 1b-3 provide[s] the distribution of lysine in the TV-markers described in Table 1. The amino acid position is defined by the X-axis. The presence of a lysine residue is

Applicants: Alexander Gad and Dora Lis

Serial No.: 09/816,989 Filed: March 23, 2001

Exhibit 1

indicated by a vertical bar at the indicated amino acid position.

Page 8, paragraph 1, line 1-3

Figures 1c-1, 1c-2 and 1c-3 provide[s] the distribution of glutamic acid in the TV-markers described in Table 1. The amino acid position is defined by the X-axis. The presence of a glutamic acid residue is indicated by a vertical bar at the indicated amino acid position.

Page 8, paragraph 2, lines 4-6

Figures 1d-1, 1d-2 and 1d-3 provide[s] the distribution of tyrosine in the TV-markers described in Table 1. The amino acid position is defined by the X-axis. The presence of a tyrosine residue is indicated by a vertical bar at the indicated amino acid position.

Page 23, paragraph 1, lines 32-36

Figures 1a-1, 1a-2, 1a-3, 1b-1, 1b-2, 1b-3, 1c-1, 1c-2, 1c-3, and 1d-1, 1d-2 and 1d-3 provide the distribution of alanine, lysine, glutamic acid and tyrosine, respectively, in the TV-markers described in Table 2. The amino acid position is defined by the X-axis, with the first amino acid corresponding to the C-terminal position. The presence of an amino acid is indicated by a vertical bar at the indicated amino acid position.